

Women and Risk of HIV/AIDS Infection

Warning: DOMDocument::loadHTML(): htmlParseEntityRef: expecting ';' in Entity, line: 14 in /home/gender/public_html/plugins/content/image/emogrifier.php on line 63 Warning: DOMDocument::loadHTML(): htmlParseEntityRef: expecting ';' in Entity, line: 14 in /home/gender/public_html/plugins/content/image/emogrifier.php on line 63 Warning: DOMDocument::loadHTML(): htmlParseEntityRef: expecting ';' in Entity, line: 14 in /home/gender/public_html/plugins/content/image/emogrifier.php on line 63

AUTHOR: International Women's Health Coalition (IWHC)

DATE: 2008

PUBLISHER: International Women's Health Coalition (IWHC)

This fact sheet reviews women's vulnerability to HIV/AIDS, focusing particularly on young women. Many women and young people do not have access to the information and services they need to protect themselves from HIV infection. Women's vulnerability is increased by marriage, sex work, injecting drug use, being the partner of an injecting drug user, a student, and violent or coercive sex.

Article can be accessed on-line [here](#) in PDF format.

To view PDF documents you will need to download and install Adobe Acrobat Reader. The program is available for free through the Adobe website, download [here](#). More help is available on PDF documents and Adobe Acrobat Reader [here](#)